



Response of laying Japanese Quails (*Cotunix japonica*) to Sandbox (*Hura crepitans*) seed meal-based diets

Nsa, E. E¹, Umoren, E. P^{1*} , Archibong, E. E¹ and Ukoha, O. A²

¹Department of Animal Science, University of Calabar, Calabar, Cross River State, Nigeria.

²Department of Animal Nutrition and Forage Science, College of Animal Science and Animal Production, Michael Okpara University of Agriculture, Umudike, Abia State, Nigeria.

*Corresponding Author
Umoren, E. P

Department of Animal
Science, University of
Calabar, Calabar,
Cross River State,
Nigeria.

Article History

Received: 01.06.2025
Accepted: 04.07.2025
Published: 16 .07.2025

Abstract: A 70-day study was carried out to assess the effects of sandbox (*Hura crepitans*) seed meal-based diets on haematology and egg production parameters of Japanese quail birds. A total of 200 seven weeks old female Japanese quails were selected from growing phase and randomly distributed to five dietary treatments with four replicates of 10 birds each in a completely randomized design. The diets were formulated to contain 0, 25, 50, 75 and 100% of sandbox seed meal (SBSM) at the expense of soybean meal to represent treatment diets 1, 2, 3, 4 and 5 respectively. Fresh drinking water was provided *ad libitum* throughout the laying phase. Blood samples were collected from two birds in each replicate at the end of the study for haematological studies. The results showed that hen day production, peak production, age at peak production and feed cost per dozen (₦) were significantly ($p < 0.05$) affected. Hen day production decreased at 100% sandbox seed meal replacement in the diets while, peak production decreased at 75 and 100% replacement level. Egg quality parameters were not significantly affected by treatment diets except the egg weight and yolk weight which had increased values with dietary increase in sandbox seed meal. Apart from packed cell volume and haemoglobin which showed significant ($p < 0.05$) differences, only at T5 (100% SBSM), other blood parameters indicated no significant ($p > 0.05$) differences across the treatment means. It could be concluded that sandbox seed meal can replace up to 75% of soya bean meal in Japanese laying quails without sacrificing egg quality, quantity indices and the health status of the birds.

Keywords: *Hura crepitans*, Japanese quails, Egg parameters, Haematology, Egg production.

Cite this article:

Nsa, E. E., Umoren, E. P., Archibong, E. E., and Ukoha, O. A., (2025). Response of laying Japanese Quails (*Cotunix japonica*) to Sandbox (*Hura crepitans*) seed meal-based diets. *ISAR Journal of Multidisciplinary Research and Studies*, 3(7), 10-17.

Introduction

Despite the great potential of quail production in Nigeria, uncontrolled increase in prices of conventional feed ingredients such as maize, soya bean and groundnut meal which serve as the principal sources of energy and protein has skyrocketed the cost of production of quail birds (Kolo and Charles, 2019; Mnisi et al., 2021). Soya bean accounts for about 25-35% of poultry feed (Adli, 2021; Tommer et al., 2024), therefore replacing soya bean in poultry feed with less costly feedstuffs will significantly reduce cost of animal products (Edacheet et al., 2007). The search for locally available, unconventional, low cost but nutritionally adequate feedstuff has therefore become imperative (Oke et al., 2025). One of such is sandbox (*Hura crepitans*) seed meal.

Sandbox tree is 90-130 feet high tropical evergreen tree of the spurge family *Euphorbiaceae*, native to tropical regions of

North and South America including the Amazon Rainforest (Borokini and Omotay, 2012). Apart from the name sandbox tree, it has several other names like Possum wood, monkey's dinner bell, monkey's pistol and Monkey no-climb (Owojuyibe et al, 2020). It grows on clay soil in moist areas near water courses and in flooded areas (Oraegbunam, et al 2020). Sandbox tree has been introduced in California, Florida, the Bahamas and various Tropical Countries around the world including Nigeria (Oraegbunam et al., 2020).

The seeds are rich in minerals like potassium (238.33mg/100g) and sodium (71.67mg). Both minerals play a vital role in active transport across the cell membrane and they are required for maintenance of osmotic balance (Hassan et al 2018; Mohammed et al., 2013). It is also rich in Magnesium and copper. The former activates enzymatic system responsible for calcium metabolism in the bone and nerves while the latter plays a role in



haemoglobin formation and contributes to iron and energy metabolism (Hassan et al., 2018).

Hura crepitans seeds also contain oil which enhances the energy density of a diet for normal maintenance and productive functions. It also serves as a source of essential fatty acids as well as carrier of the fat-soluble vitamins (Nsa et al., 2025). The dry matter of the seed (91-95%) is quite comparable to those of the conventional feedstuff, an indicator of having a long store life devoid of mold growth. The seed is moderate in crude protein 30.22% (Archibong et al., 2023), high in ether extract of about 38.95-51.24% (Yaakugh et al., 2001) over 100 times the value for the conventional oil seed. A real good source of oil and energy in livestock feed (Ansari-poor et al., 2020). However, the seed is known to contain some anti-nutritional factors like alkaloids (5.00mg/100g), tannins (5.11mg/100g), saponin (2.2mg/100g and trypsin inhibitor (30.27TIU/mg) however, there are all heat labile, which means if well treated could be a good source of both protein and energy in livestock feeds.

This research therefore was aimed to assess the response of laying Japanese quails (*Coturnix japonica*) to different replacement levels of soya bean seed meal with sandbox (*Hura crepitans*) seed meal.

Materials and Methods

Experimental diets and feeding

Five experimental layer's diets (Table 1) were formulated to contain (0, 25, 50, 75 and 100%) of sandbox seed meal, at the expense of soya bean meal, to represent treatments 1, 2, 3, 4 and 5, respectively. The diet provided the required nutrients of laying Japanese quails ration according to NRC (1994) recommendations. The quails were fed a known quantity of feed between 07:00 - 08:00am and also in the evening 16:00 - 17:00pm hours daily. Leftover feed was measured the next morning.

Table 1: Gross composition of Japanese quail sandbox seed meal-based layer's diets.

Ingredients	Replacement levels of soya bean meal with sandbox seed meal				
	T1 (0%)	T2 (25%)	T3 (50%)	T4 (75%)	T5 (100%)
Maize	42.40	42.40	42.40	42.40	42.40
SBSM	0.00	8.00	16.00	24.00	32.00
Wheat offal	7.00	7.00	7.00	7.00	7.00
Palm kernel cake	5.80	5.80	5.80	5.80	5.80
Soybean meal	32.00	24.00	16.00	8.00	0.00
Fish meal	2.50	2.50	2.50	2.50	2.50
Bone meal	5.50	5.50	5.50	5.50	5.50
Oyster shell	4.00	4.00	4.00	4.00	4.00
Salt	0.20	0.20	0.20	0.20	0.20
Lysine	0.20	0.20	0.20	0.20	0.20
Methionine	0.20	0.20	0.20	0.20	0.20
*Vit./Min. Premixes	0.20	0.20	0.20	0.20	0.20
Total	100.00	100.00	100.00	100.00	100.00
Calculated composition %					
Crude protein	24.60	24.10	23.76	23.42	23.08
Crude fiber	5.13	5.34	5.92	6.50	6.98
ME (Kcal/Kg)	2505.10	2516.18	2527.26	2538.34	2549.42
Analyzed Values %DM					
Crude protein	24.79	24.63	24.46	24.21	24.08
Crude fibre	5.11	5.34	5.49	5.97	6.23
Ether extract	3.36	3.92	4.50	4.98	5.23
ME(Kcal/kg)	2747.22	2750.87	2815.76	2828.43	2831.92
Calcium	3.32	3.26	3.24	3.23	3.30
Phosphorus	1.79	1.79	1.73	1.71	1.70

*To provide the following per kg of feed, vit A:3,400.00 i.u, vit D: 600,000.00 Iu., vit E: 4,000.00mg; vit K: 400.00mg, niacin: 6,000.00mg, vit B2; 1,800.00mg, vit B6 1,200.00mg, vit B12: 600mg, pantothenic acid: 1800.00mg, choline chloride: 70,000.00mg, cobalt: 80,000.00mg, copper: 1200.00mg, iodine: 400.00mg, iron; 8,000.00mg, manganese; 16,000.00mg, selenium: 80.00mg, zinc; 12,000.00mg

SBSM= Sandbox seed meal

Data collection

During the laying phase, the initial weight of quails were taken at the start of the experiment and thereafter on a weekly basis. Weight gain was obtained by subtracting the initial weight from the final weight. Feed intake per group was determined by weighing the feed offered in grams and the left over in grams the next morning and finding the difference. These we used to calculate the following;

$$\text{Feed intake/bird/day} = \frac{\text{Quantity of feed given(g)-leftover feed}}{\text{No of birds} \times 70 \text{ days}}$$

$$\text{Weight gain/bird/day} = \frac{\text{Final live weight-initial weight}}{\text{No of birds} \times 70 \text{ days}}$$

Protein intake

$$= \frac{\text{CP content of the diet} \times \text{quantity of feed consumed}}{100}$$

$$\text{Protein efficiency} = \frac{\text{Weight gain}}{\text{Protein intake}}$$

Feed conversion ratio (kg of feed per dozen egg)

Egg production was recorded daily and pooled weekly to calculate the hen-day production. Cost per dozens of eggs was calculated based on the prevailing market prices of the feed ingredients. All eggs laid each day in a particular replicate group were weighed to determine average egg weight.

Eggs were collected daily and weighed throughout the 70 days of the laying phase, they were randomly selected per replicate for external and internal egg quality evaluation by methods described by Tamiru et al. (2019).

Egg quality parameters

Two freshly laid eggs free of cracks were randomly selected from each replicate for weekly measurement of egg quality. Each egg was assessed separately for external and internal egg quality traits. External egg parameters; these included; egg weight (measured by the use of Hana weighing scale), egg length (using Vernier sliding caliper) and egg diameter which was measured at three points; on the blunt end, equatorial region and at the pointed end using the same vernier sliding caliper.

Internal egg quality; these were determined using the following parameters; Albumen index (determined by using micrometer screw gauge to obtain the height and weight of the albumen), yolk index (obtained as ratio and height to the width of the yolk using vernier sliding caliper), yolk colour (obtained by matching the color of the yolk with Roche yolk color fan), Haugh unit; measured by the use of spherometer screw gauge and applying the formula;

$$100 \log (H+7.57-1.7W0.37)$$

Where, H is the observed albumen height in millimeter

W is the observed weight of egg in gram (Haugh, 1937), and shell thickness (measured with a micrometer screw gauge).

Cost of production

Cost analysis using appropriate formulae were used to determine the economics of production based on the prevailing local market price of feedstuff.

$$\text{Feed cost/kg} = \frac{\text{Sum of prices of ingredients}}{100\text{kg}}$$

$$\text{Total feed cost} = \text{Total feed intake} \times \text{feed cost/kg}$$

$$\text{Total cost of feed consumed/kg (N)} = \text{Cost of feed/kg (N)} \times \text{total feed consumed (kg/bird)}$$

$$\begin{aligned} \text{Cost of feed/kg live weight gain (N)} \\ = \frac{\text{Total cost of feed consumed/kg (N)}}{\text{Total weight gain (g/bird)}} \end{aligned}$$

Haematological analysis

At the end of the feeding trial, two quails per treatment were selected for blood analysis. Blood samples were collected from the veins of the wings web with the use of hypodermic syringe into a clean test tube containing anticoagulant-Ethylene Diamine Tetra Acetate (EDTA) and used for the determination of haematological parameters. The concentration of red blood cells, white blood cells, packed cell value and haemoglobin were evaluated as described by Oriade et al. (2021). Other haematological indices including mean corpuscular haemoglobin concentration (MCH), Mean corpuscular volume (MCV), mean corpuscular haemoglobin concentration (MCHC), platelet and lymphocytes, were determined using appropriate formulae as described by Akintunde et al., 2017.

Statistical analysis

Data collected were analyzed using one-way analysis of variance of (SAS, 2010) statistically package and the significant differences between means were compared at 5% confidence level using Duncan multiple range test.

Results and Discussion

Egg production parameter of quail birds fed sandbox seed meal (SBSM) based diets are represented in table 2. The eggs laid per quail ranged from 47.00- 59.00g and showed no significant ($p>0.05$) differences across the various treatments. The dozen egg per quail ranges from 3.92-4.92g of birds in the various treatments with significant decrease ($p<0.05$) only at 100% replacement level of soya bean meal with sand box seed meal. This observation could imply that quail birds can be fed with sandbox seed-based diets up to 75% level of replacement conveniently without having negative effect on egg produced. This is in line with findings by Ozeudu et al. (2020), who worked with layer chicken with the same test ingredients but observed a drop in egg production, as the level of soya bean replacement with sand box seed meal increases above 25%. Quail birds are hardy birds, can tolerate more fibrous and less nutritious feeds (Sogunle et al., 2007). Sand box seed meal is of higher fibre content than soya bean meal as shown in table 1, this could explain why the drop was only after 75% replacement level of soybean meal.

The mean daily feed intake of birds ranged between 27.09-29.02g and did not differ ($p<0.05$) significantly among the dietary

treatments. However, there was an observed lower value recorded for birds on T5. This observation is in tandem with the report of Ukachukwu and Akpan (2007), who observed significant lower feed intake of layer birds on fibrous feedstuff than the control diet. Sandbox seed meal had crude fibre level of (12%) relatively lower than soya bean meal. Therefore, increment of the sandbox seed meal in the diet, will definitely lead to increase in the fibre level of the diet. Sandbox seed meal has higher oil content than soya bean meal (Table 1), hence high energy density as the meal increases in a diet, which is in line with separate findings of Odunsi and Gbadamosi (2001); Agida et al. (2020) that monogastric animals are expected to consume less of high energy diet in attempt not to overshoot their energy requirements. An earlier report by Aduku and Nuhu (1996) also supported this view on broiler birds fed diets containing oil sludge, which showed a slight but non-significant ($p>0.05$) decrease in feed intake as the dietary oil level increased. In support of this, Longe and Adekoya (1998) also reported that addition of oil to a diet that is otherwise adequate in energy, lower feed intake.

The mean feed conversion ratio (feed/dozen egg) was not significantly ($p>0.05$) affected by the dietary treatments. However, the hen-day production was significantly ($p<0.05$) affected by dietary treatments with birds fed diet 100% replacement of soya bean meal with sandbox seed meal having the least HDP, while other treatment diets including the control had statistically similar but higher ($p>0.05$) values.

The hen-day production (HDP) of the birds on the control groups and birds on diet 2, 3 and 4 were significantly ($p<0.05$)

higher than birds on T5. This observation supports the reports by Odetola et al. (2017) for Japanese quail birds fed kenaf seed meal. This means that replacing soybean meal with sandbox seed meal in laying Japanese birds up to 75% replacement with soya bean meal have no effect on hen-day production.

The peak egg production showed significant ($p<0.05$) differences, with birds on the control (T1), T2 and T3 having numerically ($p>0.05$) similar values that were significantly ($p<0.05$) higher than T4 and T5.

Age at peak of production showed significant ($p<0.05$) differences. Birds on the control group and T3 diets reached peak of production earlier than birds on other diets. It took birds on treatment 5, 10 days later when compared with the control to reach their peak of production.

Mortality percentage across the treatments were low and showed no significant ($p>0.05$) difference. An indication that sandbox seed meal is not lethal to Japanese birds.

Cost per dozen egg production (N) showed significant ($p<0.05$) differences. Birds on the control and 100% SBSM had similar ($p>0.05$) values that were higher than T2. Birds on T3 and T4 had the least cost per dozen egg production (N)

The observed reduction of cost per dozens of eggs produced for birds on sandbox-based diets apart from birds on T5 could be attributed to the high cost of soya bean meal and the corresponding number of eggs laid.

Table 2: Egg production parameters of Japanese quail birds fed sandbox seed meal-based diets.

Parameters	Percentage inclusion level of sandbox seed meal					SEM
	T1 (0%)	T2 (25%)	T3 (50%)	T4 (75%)	T5 (100%)	
Egg laid/bird(g/b)	59.00	58.00	53.00	50.00	47.00	1.88
Dozen egg/bird(g/b)	4.92	4.83	4.42	4.17	3.92	0.06
Feed/hen/day(g/b/d)	28.88	28.41	29.02	27.82	27.09	0.91
FCR;feed/egg(g)	4.33	4.21	4.19	4.14	5.11	0.30
Hen-day production (%)	76.00 ^a	75.77 ^a	72.00 ^{ab}	70.65 ^{ab}	65.43 ^b	3.11
Peak egg production (%)	85.00 ^a	84.99 ^a	81.57 ^a	76.67 ^{ab}	70.43 ^b	7.09
Feed cost/bird(₦)	8.58 ^a	8.16 ^b	8.01 ^b	7.38 ^c	6.86 ^d	0.71
Feed cost per dozen egg(₦)	265.45 ^a	233.32 ^b	219.40 ^c	218.90 ^c	269.51 ^a	6.98
Age at peak production day	128 ^b	130 ^b	134 ^{ab}	134 ^{ab}	141 ^a	3.99
% Mortality	2.25	1.50	1.65	2.65	3.00	0.32

^{abc}Means within the same row with different superscripts are significantly different from one another ($p<0.05$)

Egg quality parameters of quail birds fed dietary levels of SBSM are represented in Table 3. There were no significant ($p>0.05$) differences in the external and internal egg quality parameters except on the egg weight, yolk weight and yolk color. The non-significant ($p>0.05$) Differences in values of egg quality parameters were expected since the birds were of the weight and mature at the same age. This observation supports the findings of

Abdullah et al. (2003); Perez-Bomilla et al. (2012) and Zita et al. (2013) who reported that age of the hen strongly influenced the egg size and the proportion of its components. The egg yolk weights tended to be significantly ($p<0.05$) higher as the level of sandbox seed meal increased in the diets. There was a significant ($p<0.006$) increase at T3, T4 and T5. The observed improvement in the yolk weight as sandbox seed meal increased in the diet could be a result

of high content of oil in the diet, as contributed by sandbox seed meal which is known to be of higher oil content than soya bean meal (Table 1). This observation is in consonance with that of Barzegar et al. (2019) who worked on the effect of increasing metabolizable energy of pearl miller of corn and found out that higher metabolizable energy in diets of laying hens increased egg yolk weight.

Egg yolk is mainly made of carbohydrate, oil and protein (Adedeji et al., 2008), and the egg height, shell weight, yolk height, yolk diameter, yolk index, albumen weight and albumen height showed no significant ($p>0.05$) difference across dietary treatments. The shell thickness was not also affected ($p>0.05$) by dietary treatments. This is in line with research by Ozeudu et al. (2020) who reported a non-significant ($p>0.05$) difference in shell thickness of layer birds fed cooked sandbox seed meal. Shell thickness is a good indicator of shell strength and is important in the handling and marketing of eggs to minimize breakage. It

therefore appears that sandbox seed is not lacking in essential minerals and vitamins which are directly involved in formation of strong egg shell.

The yolk color became less yellowish as the level of sandbox seed meal inclusion increased in the diets, an indication of reduced presence of Vitamin A precursor in Sandbox seed meal. Haugh unit ranged from 79.95 – 84.44 in line with the report of Ukachukwu and Akpan (2007) and showed no significant ($p>0.05$) differences across the treatment means and indicate high quality eggs.

Haugh unit is the major parameter for determining egg quality. According to United State of America Department of Agriculture (USDA), a Haugh unit score of 72 and above (Score AA) is acceptable and connotes freshness in egg. Quan et al. (2021) reported Haugh unit range of 76.90 to 78.24 and 78.98 to 79.10, respectively for layer birds.

Table 3. Egg quality parameters of Japanese quail birds fed sandbox seed meal-based diets.

Parameters	Replacement levels of soybean meal with sandbox seed meal					SEM
	T ₁ (0%)	T ₂ (25%)	T ₃ (50%)	T ₄ (75%)	T ₅ (100%)	
Egg weight(g)	10.23 ^b	10.26 ^b	10.33 ^{ab}	10.39 ^{ab}	10.42 ^a	0.88
Egg height(mm)	30.90	30.92	30.89	30.90	30.90	0.45
Shell weight(g)	0.81	0.81	0.80	0.80	0.81	0.01
Yolk weight(g)	3.11 ^b	3.11 ^b	3.15 ^{ab}	3.17 ^a	3.18 ^a	0.21
Yolk height(mm)	1.38	1.37	1.38	1.38	1.38	0.02
Yolk diameter(mm)	2.09	2.09	2.08	2.08	2.06	0.09
Yolk index	0.51	0.57	0.54	0.63	0.59	0.01
Yolk color	2.64 ^a	2.54 ^b	2.50 ^b	2.21 ^c	2.00 ^d	0.02
Albumen weight(g)	6.47	6.47	6.50	6.50	6.51	0.47
Albumen Height(mm)	6.45	6.49	6.50	6.50	6.58	0.36
Shell thickness (mm)	0.26	0.26	0.25	0.25	0.25	0.01
Haugh unit	84.44	81.31	81.12	80.79	79.95	5.66

^{abcd} means within the same row with different superscripts are significantly different from one another ($p<0.05$)

The haemoglobin values obtained in this study ranged from 13.17g/dl to 20.09g/dl. These were higher than the normal range for chickens (7-13g/dl) and the range (8.7 to 9.3g/dl) as reported by others (Atansuyi et al., 2019), for indigenous chickens, but within the range reported by Agina et al.(2017) for healthy quail birds. The observed differences in Hb values could be attributed to differences in bird type and test ingredients used.

Eggun (1989) reported that, high or low Hb could be attributed to a state of nutritional anaemia in birds and the quality of protein fed to the birds. Therefore, the observed lower Hb obtained by birds on diet T5 was in line with the report of (Nsa et al., 2025). that Hb levels were positively correlated with protein quality than sandbox seed meal, this is reflected in the Hb levels of

the different diets. Increase in the haemoglobin level may be accompanied by a rise in the red blood cells and packed cell volume (Jain et al., 2011)

The RBC, MCV, MCH and MCHC values obtained in this study were within the ranges reported by (Atansuyi et al. (2019) for a healthy chicken. This shows that, the bone marrow of the birds were functioning normally, also the normal values obtained for MCV and MCHC in this study indicated that, there was no negative interaction between energy and protein levels in the diets (Akinola and Egwuanumku (2017). A reduction in MCV occurs when iron deficiency becomes severe, which triggers anaemia. This was not a case here as its level was within a normal range for poultry birds.

Table 4. Haematological indices of laying Japanese quail birds fed sandbox seed meal-based diets.

Parameters	T1 (0%)	T2 (25%)	T3 (50%)	T4 (75%)	T5 (100%)	SEM	Ref. Ranges <i>Agina et al. (2017)</i>
PCV (%)	56.00a	54.08a	53.11a	49.89a	30.67b	1.04	25-66
RBC($\times 10^3/L$)	4.03	4.00	3.96	3.93	3.90	0.05	2.02-6.99
Hb(g/dl)	20.09a	20.00a	19.87a	19.22a	13.17b	0.65	8.61-20.57
WBC($\times 10^6/L$)	120.22	119.02	117.41	121.00	118.65	10.05	100.00-125.00
MCV(fl)	118.20	120.11	122.03	130.31	130.66	8.62	64.68-131.99
MCH(g/dl)	48.72	49.38	59.62	60.54	60.73	4.11	20.72-69.08
MCHC(g/dl)	39.09	39.21	40.55	45.29	47.66	1.54	20.77-64.28
Lymphocytes	56.20	56.90	54.12	70.71	60.40	9.98	30.05-88.50
Platelet ($\times 10^6/L$)	30.21	35.02	32.28	36.41	36.82	5.81	22.65-45.60

^{ab} Means within the same row with different superscript are significantly different ($p < 0.05$)

Conclusion

This study showed that up to 75% of soybean meal could be replaced with Sandbox seed meal in laying Japanese quails' diet without any adverse effect on their haematological indices and egg production parameters.

References

- Abdullah, A. R., Sokunbi, O. A., Omisola, O. O. and Adewusi, M. K. (2003). Interrelations between body weight and body linear measurements in domestic rabbits (*Oryctolagus cuniculus*). *Proceedings of the 28th Annual Conference of Nigerian Society for Animal Production*. Pp. 133-136
- Adindu, F. A., Elekwa, I and Ogwo, J. J. (2016). Phytochemical comparative screening of aqueous extracts of the leaves, stem barks and roots of *Hura crepitans* (L) using GC-FID. *IOSR Journal of Biotechnology and Biochemistry*, 2, 2455-2463.
- Aduku, A. O. and Nuhu, A. (1996). A comparative evaluation of cotton seed meal oil sludge and palm oil in broiler in broiler diets. *Tropical Agriculture (Trinidad)*, 73(3):206-210.
- Adedeji, O. S., Farinu, G. O., Olayemi, T. B., Ameen, S. A. and Babatunde, G. M. (2008). Performance and egg quality parameters of laying hens fed different dietary inclusion levels of bitter kola (*Garcina kola*). *Resource Journal of Poultry Science*, 2 (4), 75-77.
- Adli, D. N. (2021). Use of insects in poultry feed as replacement soya bean meal and fish meal in development countries: a systematic review. *Livestock Research for Rural Development*, 33, 128. <http://www.lrrd.org/lrrd33/10/33128danun.html>
- Agida, C. A., Amaduruonye, W., Nsa, E. E. and Nathaniel, J. (2020). Serum biochemistry, haematological profile and organ proportion of broiler starter chicks fed graded levels of palm oil mill effluent (POME). *Journal of Animal Science and Veterinary Medicine*, 5(1), 202-211.
- Agina, O. A., Ezema, W. S. and Nwishiényi, C. N. (2017). Haemato-biochemical profile of apparently healthy domestic adult Japanese Quails in Nsukka, Enugu State Nigeria. *Animal Research International*, 12(1), 2120-2129.
- Akinola, L. A. F. and Egwuanumku, J.O. (2017). Haematology and serum biochemistry of laying hens fed red pepper (*Capsicum annum* L.) as feed additive. *Nigerian Journal of Animal Science*, (1), 127 - 134
- Akintunde, A. R., Omege, J. J., Bawa, G. S., Onimisi, P. A. and Samuel, I. (2017). Haematological parameters and serum biochemistry of Japanese quail chicks (*Coturnixcoturnix japonica*) fed raw and processed pigeon pea (*Cajanus cajan*) seed meal based diets. *Nigerian Journal of Animal Production*, 44(1), 89 - 96.
- Ansariipoor, A., Sedghi, M., Sadeghi-Sefidmazgi, A. and Pilevar, M. (2020). Optimization of growth performance responses of Japanese quail with different concentrations of metabolizable energy, lysine, and sulfur amino acids using Taguchi method. *Livestock Science*, 241. <https://doi.org/10.1016/j.livsci.2020.104234>.
- Archibong, E. E., Nsa, E. E. and Umoren, U. E. (2018). Nutritional evaluation of tigernut (*Cyperus esculentus*) meal as a replacement for maize in broiler diets. *Nigerian Journal of Animal Production*, 45 (4), 90-98.
- Atansuyi, A..J., Ogunribido, T. Z. and Chineke, C. A. (2019). Haematological and serum biochemical characteristics of four chicken genotypes in south-western, Nigeria *Nigerian J. Anim. Sci.* 21 (1), 9-16.
- Barzegar, S., Wu, S., Noblet, J., Swick, R. A. (2019). Metabolizable energy of corn, soybean meal and wheat for laying hens. *Poultry Science*, 98 (11), 5876-5882. <https://doi.org/10.3382/ps/pez333>.
- Borokini, T. I. and Omotayo, F.O. (2012). Phytochemical and ethnobotanical study of some selected medicinal plants from Nigeria. *Journal of Medicinal Plant Research*, 6(7), 1106-1118.

15. Edachet, J. A., Musa, U., Karsin, P. D., Esilonu, J. O., Yisa, A., Okpala, E. J. and Zwandor, N. J. (2007). Evaluation of the performance of growing Japanese Quails (*Coturnix coturnix japonica*) fed graded levels of banana peel meal. In: Sustainability of the livestock industry in an oil Economy. (Editor, Agiang, E.A., Agwunobi, L.N. and Olawoyin, O. O.). *Proceedings of the 32rd annual conference of the Nigerian Society for Animal Production*, Calabar, March, 18-21, 2007, 342-346.
16. Eggun, B.O. (2009). Blood urea measurements as a technique for assessing protein quality. *British Journal of Nutrition*, 24, 983-988.
17. Esonu, B. O., Ozeudu, E., Emenalom, O.O., Nnaji, C. and Onyeikegbulem, I. K. (2014). Nutritional value of sandbox (*Hura crepitans*) seed meal for broiler finisher birds. *Journal of Natural Sciences Research*, 4(23); 89-99.
18. Fafiolu, A. O., Oduguwa, O. O., Ikeobi, C.O.N. and Onwuka, C. F. I. (2006). Utilization of malted sorghum in the diet of rearing pullets and laying hens. *Arch. Zootechnology*, 55 (212), 361- 371.
19. Hassan, L. G. Sokoto, A. M., Ngaski, M. N., Anka, S. A., Chanchangi, B. M., Umar, K. J. and Keay, R.W.J. (2018). Trees of Nigeria. *Nordic Journal of Botany*, 11, 322.
20. Haugh, R. R, 1937. The Haugh unit for measuring egg quality. *U.S. Egg Poultry Magazine*, 43, 522-555, 572-573.
21. Jain, P. C., Kataria, N., & Kataria, A. K. (2011). Haematological studies in quail. *Journal of Veterinary and Animal Sciences*, 1(1), 1-5.
22. Kabir, A. (2013). Blood chemistry analyses of Japanese quail (*Coturnix coturnix Japonica*). *Scholarly Journal of Agricultural Science*, 3(4), 132-136.
23. Kolo, Y. S. and H Charles, H. (2019). Cost effectiveness of Japanese quails (*Coturnix coturnix japonica*) fed fermented mango (*Mangifera indica*) kernel composite meal as a replacement for maize. *International Journal of Agriculture and Biosciences*, 8(5), 222- 225.
24. Longe, O. G. and Adekoya, O. E. (1988). Response of laying hens to different inclusion levels of palm kernel meal and vegetable oil. *Nigerian Journal of Animal Production*, 15, 111-117.
25. Mnisi, C. M., Marareni, M., Manyeula, F. Madibana, M. J. (2021). A way forward for the South African quail sector as a potential contributor to food and nutrition security following the aftermath of covid-19: A review. *Agriculture and Food Security*, 10, 48. <https://doi.org/10.1186/s40066-021-00331-8>
26. Muhammed, N. A., Isiaka, A.A. and Adeniyi, O. A. (2013). Chemical Composition of *Hura crepitans* seeds and antimicrobial activities of its oil. *International Journal of Science and Research*, 2 (3), 2319-2324.
27. NRC (1994). National Research Council. Nutrient requirements of domestic animals. Nutrient requirement of poultry, 4th edition. National Academy Press Washington D.C.100-120
28. Nsa, E. E., Ozung, P. O. & Archibong, E. E. (2025). Chemical composition and feeding value of sandbox (*hura crepitans*) seed meal as an alternative plant protein source for rabbits in Southern Nigeria. *Global Journal of Pure and Applied Sciences*, 31, 559-571. <https://dx.doi.org/10.4314/gjpas.v31i3.12>
29. Odetola, O. M., Adejinmi, O.O., Akingbade, A. O. and Adekanmbi, A. O. (2017). Performance and egg quality parameters of Japanese quail (*Coturnix coturnix japonica*) fed kenaf (*Hibiscus cannabinis* L.) seed meal-based diets subjected to varying cooking periods. *Nigerian Journal of Animal Production*, 44(4), 225-231.
30. Odunsi, A.A. & Gbadamosi, A.J. (2001). Effects of dietary inclusion of palm oil and shea butter fat on growth and sexual maturity of pullets. *Nigerian Journal of Animal Production*, 28(1):26-30.
31. Oke, O. E., Oliyide, K. M., Akosile, O. A., Oni, A. I., Adekunle, E. O., Oyebanji, B. O., Aremu, O. P., Adeoba I. M., Eletu, T. A. & Daramola, J. O. (2025). Innovations in quail welfare: Integrating environmental enrichment, nutrition and genetic advances for improved health and productivity. *Veterinary Medicine and Science*, 11, <https://doi.org/10.1002/vms3.70424>
32. Oraegbunam, J. C., Oladipo, B., Falowo, O. A., and Betiku, E. (2020). Clean sandbox (*Hura crepitans*) oil methyl esters synthesis: A kinetic and thermodynamic study through pH monitoring approach.. *Renewable Energy*, 160, 526-537, <https://doi.org/10.1016/j.renene.2020.06.124>.
33. Oriade, W. A., Omoleye, O.S., Adebayo, F. B. and Adigun, O.T. (2021). Haematological parameters of Japanese Quail (*Coturnix Coturnix Japonica*) raised on deep litter system in South Western Nigeria. *Elixir Agriculture*, 161, 55804-55807
34. Owojuyigbe, O. S., Firemp, C., Laribie, C. and Komlaga, K. (2020). A Review of Ethnomedical Phytochemical and Pharmacological Studies. *Journal of Complementary and Integrative Medicine*, 9(2), 1-10.
35. Ozeudu, E., Esonu, B.O., Udebibie, A.B.I., Ojike, F. C., Uchegbu, M. C., Emenalom, O. O. and Chibuike, C. E. (2020). Substitution value of cooked sandbox (*Hura crepitans*) seed meal in layers diets. *Nigerian Journal of Animal production*, 42(1), 79-84.
36. Perez-Bomilla, A. S., Novoa, J., Garcia, M., Mohifi-Ashi, M. and Meteos, G. G. (2012). Effects of energy concentration of the diets on the productive performance and egg quality of brown egg lay hens differing in initial body weight. *Poultry Science*, 91, 3156-3166.
37. Quan, C., Qian Xi, Q., Shi, X., Han, R., Du, Q., Forghani, F., Xue C., Zhang, J. and Wang, J., (2021). Development of predictive models for egg freshness and shelf-life under different storage temperatures, *Food Quality and Safety*, 5, 1-7. <https://doi.org/10.1093/fqsafe/fyab021>
38. SAS Institute, 2010. SAS Proprietary Software, Release 9.2 Cary, NC, SAS Inst. Inc.

39. Sogunle, O. M, Fanimu, A. O., Abiola, S.S. and Bamgbose, A. M. (2007). Growth response, body temperature and blood constituents of pullet chicks fed cassava peel meal supplemented with cashew nut reject meal. *Nigerian Journal of Animal Production*, 34(1), 32-44.
40. Tamiru, H., Duguma, M. , Furgasa, W. and Lema Yimer, L. (2019). Review on chicken egg quality determination, grading and affecting factors. *Asian Journal of Medical Science Research & Review*, 01(01), 34-42. <http://ajmsrr.com/index.php/ajmsrr>
41. Toomer, O. T., Oviedo-Rondón, E. O., Ali, M., Joseph, M., Vu, T., Fallen, B., & Mian, R. (2024). Full-fat Soybean meals as an alternative poultry feed ingredient—feed processing methods and utilization: Review and perspective. *Animals*, 14(16), 2366. <https://doi.org/10.3390/ani14162366>
42. Ukachukwu, S. N. and Akpan, U. O. (2007). Influence of level and duration of quantitative feed restriction on post-restriction egg-laying characteristics and egg quality of pullets. *International Journal of Poultry Science*, 6, 567-572. doi: 10.3923/ijps.2007.567.572
43. Yaakugh, I.D.I., Tuleum, C.D. & Kaankuka, F.G. (2001). MSc. Thesis. Nutrient Composition of Hura crepitans. Department of Animal Production, University of Agric, Makurdi, Nigeria.Pp 87.
44. Zita, L., Z. Ledvinka, L. Klesalová Ita, L., Z. Ledvinka, L. Klesalová (2013). The effect of the age of Japanese quails he effect of the age of Japanese quails on certain egg quality traits and their relationships. *Veterinarski Arhiv*, 83, 223-232.